

# **BRAIN AND SPINAL CORD INJURIES**

## **Abstract**

Brain and spinal cord injuries are common medical emergencies. Brain injuries associated with medical conditions are the third leading cause of death in the United States, and they are the leading cause of disability. If there is an injury to the brain or the spinal cord, this process is disrupted. This disruption can lead to devastating consequences on bodily functions. Trauma or a medical condition may cause injury to the brain or spinal cord. The most common type of brain injury is a stroke, which is also a leading cause of disability and death in the United States. Treatment and care of a patient with a brain or spinal cord injury depends on the cause of the injury, and requires attention to the patient's nutrition, hydration, and the prevention of complications related to skin care and aspiration.

## **Learning Objectives:**

1. Discuss the general functions of the brain and spinal cord.
2. Identify the most common causes of brain and spinal cord injuries.
3. Discuss a Certified Nursing Assistant's role in caring for a person with a brain or spinal cord injury.

## **Introduction**

The brain, the spinal cord and the peripheral nerves comprise the nervous system. The nervous system controls a person's conscious and unconscious behavior and bodily functions. If there is an injury to the brain or the spinal cord, this process is disrupted and can lead to devastating consequences on bodily functions. Trauma or a medical condition may cause injury to the brain or spinal cord. Treatment and care of a patient with a brain or spinal cord injury depends on the cause of the injury. A Certified Nursing Assistant (CNA) caring for a patient with a brain or spinal cord injury will generally focus on aspiration prevention, adequate nutrition and good hydration, observation for complications, and skin care.

## **Prevalence of Brain and Spinal Cord Injuries**

Brain and spinal cord injuries are common medical emergencies. Brain injuries associated with medical conditions are the third leading cause of death in the United States, and they are the leading cause of disability. Most brain injuries are a complication of widespread diseases such as atherosclerosis, diabetes, and hypertension.

Brain injuries usually happen to people over the age of 65. Traumatic brain injuries cause approximately 40% of all deaths from accidents and injuries, and each year approximately 52,000 people die from a traumatic brain injury. Spinal cord injuries are caused by trauma or accidents and most of the victims are young men. Brain injuries and spinal cord injuries can be devastating. Many of the victims who suffer a brain injury or a spinal cord injury do not survive and many of those who do survive have serious, permanent disabilities.

## The Nervous System

The brain, the spinal cord and the peripheral nerves are the three major components of the nervous system. The nervous system is the part of the body that controls everything a person does. All of a person's conscious and unconscious behavior starts with, and is controlled by, the nervous system. Although the brain and spinal cord work together, they have different, special roles.

The brain could be considered the command center of the body, and it is divided into different, specialized regions. Each region is involved in a particular aspect of functioning and behavior that a person controls - speech, emotions, problem solving, and voluntary movements such as walking. There are also specialized regions that initiate and control all of the functions and behavior that are involuntary, such as body temperature, breathing, digestion, and heartbeat.

These specific regions in the brain work by sending messages or commands to the organs and the body and receiving information and feedback from tissues and organs. For example, if a person is running to catch a train, the person's muscles work harder and need more oxygen. The muscles send this information to the cardiovascular center of the brain and respiratory center of the brain. The cardiovascular center sends a message to the heart to beat faster and a message to the blood vessels to dilate. The respiratory center sends out a message to the lungs to breathe faster and deeper. These messages result in more blood with more oxygen being delivered to the muscles.

The spinal cord is a long body of nervous tissue that is located inside the spine. It starts at the base of the brain and ends at the bottom of the spine near the buttocks. The spinal cord has two functions: 1) it helps control some basic body functions, and 2) it acts as the major “power conductor” of the information that the brain sends and receives.

Along the entire length of the spinal cord are peripheral nerves. These are long, thin strands of nervous tissue that are, in a sense, like electrical wires. These peripheral nerves travel to all of the organs, such as the heart, lungs, liver, kidneys, and glands, to receptors to sense pain, heat, or cold, and to every other part of the body. Information and messages from the different areas of the brain travel through the spinal cord and the peripheral nerves to the organs and the body. Information and messages from the organs and the body travel back to the brain through the peripheral nerves and the spinal cord.

The brain controls a person’s conscious and unconscious behavior as well as voluntary and involuntary behavior. It receives messages from the external and internal environment, for example indicating it is cold outside, or that there is a big pothole in the street in front of a person; and, of hunger or that the muscles need more oxygen. In response to these messages, the brain sends out information and commands that will cause a person to start shivering and put on a coat, step around a pothole, get something to eat to appease hunger, and increase heart rate and the rate of breathing to supply more oxygen to the muscles.

Brain and spinal cord injuries can have devastating consequences. If there is an injury to the brain or the spinal cord, this process is disrupted. Depending on the location of the injury, specific aspects of a person's behavior and functioning will be affected. For example, in some cases of brain injury a person's speech will be affected but he or she will retain all of the basic voluntary and involuntary muscle functions. Injuries to other parts of the brain will affect someone's ability to swallow or to walk. If someone suffers a severe injury to specific areas of the spinal column, that person will not be able to move the arms or legs.

### **Traumatic and Acquired Brain Injuries**

There are two basic causes of brain injuries: *trauma* and *acquired conditions*. Traumatic brain injuries are less common and will only be discussed briefly. This section will focus on brain injuries caused by complications associated with an acquired condition.

Traumatic brain injuries are caused by a violent force that is applied to the head. Motor vehicle accidents, accidents related to sports, firearm injuries, and falls are common causes of traumatic brain injury. The intensity of a trauma to the head may damage the brain and cause bleeding and swelling.

Due to the brain being enclosed inside the skull, the bleeding and the swelling cause pressure on the brain tissue. If the bleeding, swelling, pressure, and the physical damage caused by the trauma are severe and cannot be treated, the brain will suffer permanent injury.

Acquired brain injuries can be caused by cancers, cardiac arrest, cardiac arrhythmias, hemorrhage, infections, and thrombosis (commonly known as a blood clot). A stroke is the most common cause of a brain injury. A stroke is caused by hemorrhage or a thrombosis, and a discussion of stroke will be the primary focus of this section. Brain injury from cancers, cardiac arrest, cardiac arrhythmias, and infections will be briefly discussed. A transient ischemic attack (TIA) is a neurological accident that is similar to stroke.

## **Stroke**

The most common type of brain injury is a stroke. Strokes are one of the leading causes of death in the United States. They are *the* leading cause of disability, and each year approximately 700,000 people in the U.S., have a stroke. The survival rate after a stroke has improved with better treatment, improved emergency care, and increased public awareness, but strokes are still a major public health problem and are considered a medical emergency.

The technical term for a stroke is *cerebrovascular accident (CVA)*. The word vascular means of or pertaining to the blood vessels and circulation. The prefix *cerebro* means *of or pertaining to the brain*. The two together provide the basic definition of a stroke or CVA.

A stroke is defined as a sudden disruption of the blood supply to the brain that may cause permanent damage. There are two types of stroke: 1) ischemic stroke and 2) hemorrhagic stroke. Ischemic strokes are more common than hemorrhagic strokes; approximately 80% of all strokes are ischemic strokes.

## Ischemic Stroke

*Ischemia* is a medical term that means decreased blood supply to a specific area. An ischemic stroke happens when the blood flow to the brain is interrupted by a blood clot.

There are two causes of ischemic stroke. The first is *atherosclerosis*, commonly called "hardening of the arteries". Atherosclerosis is a very common cardiovascular disease. Atherosclerosis causes deposits of cholesterol and fat to build up in the walls of the arteries and veins. These cholesterol and fat plaques interfere with blood flow, and they also provide a place for blood clots to form. These blood clots are called a *thrombosis*. If the cholesterol and fat plaques and the blood clots inside the arteries and veins become too large, blood flow can be severely decreased or completely stopped, and an ischemic stroke can happen.

The second cause of an ischemic stroke is an *embolism*. An embolism is a thrombosis that has broken off from an artery or a vein. The embolism moves through the circulation and becomes lodged in a blood vessel in the brain, stopping the blood flow and causing ischemic damage. Ischemic strokes are also a common complication of atrial fibrillation (a very common cardiac arrhythmia), diabetes and hypertension.

## Hemorrhagic Stroke

A hemorrhagic stroke happens when a blood vessel in the brain ruptures. This causes damage in two ways. First, the brain does not receive oxygenated blood. Second, because the brain is enclosed in the skull, the blood that leaks from a ruptured artery or vein has

nowhere to go and the buildup of blood in this enclosed space causes pressure on the brain tissue. Hemorrhagic strokes are usually a complication of chronic hypertension.

The signs and symptoms of hemorrhagic stroke are essentially the same as those of an ischemic stroke, but hemorrhagic strokes are usually more severe. People who are 45 years of age or younger who have a stroke are more likely to have a hemorrhagic stroke.

Most strokes are ischemic or hemorrhagic but a small percentage of strokes can be due to illicit drug use, migraine headaches, or a sudden spasm of a blood vessel. In some cases, the cause of the stroke cannot be determined.

Why is a stroke a medical emergency? Why can a stroke cause such serious neurological damage? The simplest answer is supply and demand. The tissues and organs of the body need oxygen to survive. Oxygen is carried by the blood to the brain, heart, kidneys, muscles, and other organs so they all receive the amount they need. Without oxygen, these organs and tissues die.

The brain is especially sensitive to a lack of oxygen. The brain accounts for only 2% of total body weight but the brain needs about 15-20% of all the blood that is pumped by the heart each minute. Brain tissue is very active and it needs a big supply of nutrients and oxygen. If the blood flow to the brain is restricted or stopped, the consequences can be severe. The brain can only tolerate very short interruptions in blood flow; after 3 minutes without oxygen serious

irreversible brain damage is likely to occur.

## Stroke Risk Factors

Strokes are more common among men and more common in African-Americans. The older someone is, the greater the risk that he or she will have a stroke, and most strokes happen to people who are 65 years of age or older.

The *controllable* risk factors for stroke are atherosclerosis, certain heart arrhythmias (especially atrial fibrillation), cigarette smoking, diabetes, hypertension, obesity, and a sedentary lifestyle. The *uncontrollable* risk factors for stroke are age, gender, ethnicity, a family history of stroke, and a previous TIA. Age increases the risk for stroke because older people have increasingly serious medical problems that are associated with stroke. It is not clear why women are more likely to suffer a stroke; it may simply be because women live longer.

The ethnic risk factor for stroke has been intensely debated. It is still not clear if the ethnic risk factor is related to ethnicity itself or health and lifestyle issues. A family history of stroke is also difficult to understand, and the contribution of genetics to stroke is not clearly understood.

Atrial fibrillation, a common cardiac arrhythmia, disrupts the normal rhythmic contractions of the atria. Instead of beating in a regular, synchronized manner, the atria beat very fast and erratically - they quiver, or fibrillate. Because of this, blood pools in the atria and clots are formed on the atrial walls. Pieces of the clots - embolisms -

occasionally break off, travel through the circulation and can occlude a blood vessel in the brain.

## Signs and Symptoms of a Stroke

The signs and symptoms of a stroke depend on what part of the brain is affected and how much damage has been done. They may include:

- Weakness in one side of the body
- Differing degrees of paralysis
- Facial droop
- Decreased level of consciousness
- Loss of consciousness
- Confusion
- Inability to speak or inability to understand speech
- Sudden vision loss
- Dizziness
- Loss of balance
- Sudden and severe headache

A quick way to determine if someone is having a stroke is to use the acronym **FAST**. Look at the victim's **F**ace to see if one side is drooping. Ask the victim to hold up his or her **A**rms and see if one or both drift down. Check and see if the victim's **S**peech is slurred. The **T** stands for **T**ime, which is critical because the sooner someone receives medical treatment the better chance that person has for recovery.

**FACE → ARMS → SPEECH → TIME**

The signs and symptoms of a stroke can be dramatic; the victim may be confused, lose consciousness, or complain of a severe headache. The signs and symptoms can be mild and subtle, as well. Someone having a stroke may simply “not feel right” and complain of dizziness, weakness, or some blurred vision. A severe stroke is usually obvious, but a mild stroke or one that is just beginning may be hard to detect. In the latter situation, an exam by a health clinician and specific diagnostic tests are needed to know if that person is having a stroke. If there is any suspicion someone is having a stroke, those observing the stroke situation should call 911. Special caution should be taken if someone has risk factors for stroke.

Some people who suffer a stroke do recover, but the recovery process is difficult. The survivors often develop complication such as bedsores, depression, pneumonia, and thrombosis; and, recovery is not a guarantee of a healthy life. Approximately 20% of all stroke victims die within the first year. Permanent disabilities are common and very serious. These permanent disabilities caused by a stroke include those listed here.

- Aphasia: Aphasia means the inability to speak. Aphasia is a common complication of stroke.
- Coma
- Paralysis: Paralysis that is caused by a stroke can affect one part of the body (such as, part of the face or one arm), or it may affect one half or one side of the body. Paralysis of one side of the body

(face, trunk, arm, and leg) is called *hemiplegia*. Paralysis affecting the lower half of the body is called *paraplegia*. Paralysis affecting the upper and the lower extremities is called *quadriplegia*.

- Visual deficits.
- Weakness: The stroke victim may not suffer from paralysis, but the stroke may cause serious muscle weakness. Muscle weakness caused by a stroke that affects one side of the body is called *hemiparesis*. Muscle weakness that affects the lower extremities is called *paraparesis*. Muscle weakness that affects the upper and lower extremities is called *quadriparesis*.

### **Treatment of a Stroke**

There are two basic aspects of treatment of a stroke: 1) immediate medical care and 2) rehabilitation. The immediate medical care of a stroke is critically important. Someone who is having a stroke needs evaluation and treatment as soon as possible. Prompt medical attention is the most important part of stroke care. If it is suspected that someone is having a stroke, 911 should be called immediately.

A patient may not realize that he or she is having a stroke; the signs and symptoms may be mistaken for another illness, or the signs and symptoms may be relatively mild and subtle. Some patients may deny the seriousness of the effects, or they may simply wait, hoping that the signs and symptoms will go away. Regardless of why, a significant number of people who are having a stroke arrive at the hospital many hours after the stroke has begun, and this is the worst possible scenario. Successful treatment of a stroke requires prompt medical care.

The reason why prompt treatment of a stroke is so important is because brain cells that are deprived of oxygen will die quickly. It should be remembered that after three minutes without oxygen, irreversible brain damage will occur. However, when someone is suffering a stroke, some parts of the brain will be injured but will not be permanently damaged if treatment is delivered quickly enough. Medical personnel who care of stroke victims often say, "Time is brain," and that is very true. The brain tissue that is dead cannot be saved, but the injured areas can be restored if treatment is delivered in a timely manner. This is especially true if the patient is having an ischemic stroke.

Ischemic strokes are treated with intravenous (IV) medications that break up the clot that is causing the stroke; these drugs are called thrombolytics. Thrombolytics are usually given in an emergency room but in some circumstances they may be given by emergency medical services (EMS) personnel on the scene or while en-route to a hospital. The goal is to deliver the thrombolytics within three hours of the onset of the signs and symptoms of an ischemic stroke; remember, *time is brain*. If the thrombolytics are given within the three hour window, injured brain tissue can be restored and the patient will be less likely to suffer permanent neurological damage.

A hemorrhagic stroke is treated with antihypertensive medication to lower blood pressure and with surgical and medical procedures that decrease the pressure on the brain. These patients need prompt treatment, as well, and for the same reason discussed in the section on ischemic stroke. Hemorrhagic strokes would not be treated with thrombolytics. Thrombolytics are blood thinners that prevent the blood

from coagulating. If they were given to someone who is bleeding from a hemorrhagic stroke the damage could be catastrophic.

Once a stroke has occurred, and after the initial medical or surgical treatment has been delivered, the patient who has had a stroke needs rehabilitation. Rehabilitation for a stroke victim is similar to the immediate medical care; it must be done promptly and the sooner it is started the better the recovery is likely to be.

The rehabilitation starts with an assessment of the patient's neurological and physical capabilities. The health clinician should determine whether the patient can speak, understand speech, and is oriented to time, place, and person. Additionally, the clinician needs to know how much physical ability the patient has, and whether the patient is paralyzed or has significant muscle weaknesses, and is able to swallow. Many patients who have had a stroke have damage to the gag reflex and they cannot eat solid food. If these disabilities are present they may, with time and rehabilitation, improve or even completely resolve.

Recovery from a stroke happens in two phases. First, there is recovery that happens in the first 3-6 months after a stroke, and this accounts for most of the recovery that is going to occur. Second, there is also recovery that happens in the following months and years.

Rehabilitation will also include monitoring for, and treating the various post-stroke complications that can occur.

It can be difficult to predict how much function a stroke victim will regain but if the damage has been severe and the patient does not

recover any function within the first few months, the outlook is poor. However, studies have shown that when the amount of damage is compared to the degree of recovery, patients who undergo extensive rehabilitation do far better than would be expected. Rehabilitation does work, and the sooner it is started the better.

## **Post-Stroke Complications**

Complications after a stroke are very common. These can involve essentially every organ system.

Aspiration can be a serious problem after a stroke. Loss of the gag reflex is common and this puts the patient at risk for aspiration and pneumonia. Dysphagia, a common complication of stroke, contributes to the risk of aspiration. Aspiration is more likely to occur if the patient is > 65 or has had a severe stroke.

Major depression is common after a stroke. The psychological burden of depression is significant but depression is also a physical risk for the patient. It is not clear why but patients who develop depression after a stroke are more likely to suffer from other complications and they are more likely to die from the stroke.

Many patients who have had stroke will be immobile for some period of time, and deep-vein thrombosis (DVT) and pulmonary embolism (PE) are likely to occur. A DVT or a PE can significantly increase the risk of death, and they can prolong a hospital stay and a patient's recovery time. Fortunately, DVT and PE can be prevented with the use of anticoagulant drugs. Immobility also puts patients at risk for contractures, skin breakdown, and pressure ulcers.

Stroke and spinal cord injuries increase the risk of developing infections such as urinary tract infections and respiratory infections. Fecal incontinence and urinary incontinence are also common complications of a stroke.

### **Spinal Cord Injuries**

Spinal cord injuries are almost always caused by a traumatic injury from an automobile accident, a fall, a firearm, or a sports accident such as a collision in football. Spinal cord injuries are less common than brain injuries caused by trauma and they are less common than ischemic and hemorrhagic strokes. Spinal cord injuries can be relatively mild and the patient may recover, or they can be severe and the patient will have a permanent disability. The signs and symptoms of a spinal cord injury are similar to those of a stroke but they tend to be more extensive and more severe. Permanent disabilities are common, and people who have had a spinal cord injury can develop the same complications that are associated with a stroke.

In general, the outlook for spinal cord injuries is not good. The life expectancy of people who survive a serious spinal cord injury is significantly decreased. Approximately 10-20% of these patients die before reaching a hospital. If a patient is completely paralyzed and does not regain any function within 72 hours of the accident, that patient will never recover voluntary movement. Many people who survive the initial accident die from complications such as pneumonia, other infections, or pulmonary emboli (blood clots in the lung).

The location of the spinal cord injury is the factor that determines how extensive a disability will be. The spinal cord is located inside the spine, and the spine is divided into four separate areas that start at the neck and extend to the buttocks: cervical, thoracic, lumbar, and sacral. The area of the spine in the neck is called the cervical spine. The most serious spinal cord injuries happen when the spinal cord in the area of the cervical spine or the thoracic spine is damaged. The closer to the skull, the worse the injury is likely to be, and damage to the spinal cord in the area of the cervical spine (commonly called C-spine injuries) often causes severe and permanent damage.

### **Treatment of Spinal Cord Injuries**

Treatment of spinal cord injuries is the same as for a stroke, which include immediate medical care and then rehabilitation. Unfortunately, spinal cord injuries cannot be treated as successfully as can a stroke. In most cases, once the spinal cord is damaged there is no way to repair it. Patients who have suffered a spinal cord injury usually have serious, permanent disabilities. If paralysis is still present 72 hours after the spinal cord injury, there is no chance for recovery. Less severe injuries may not result in permanent disabilities, but the prognosis is still poor. The complications associated with a spinal cord injury are the same as those associated with a stroke. However, because the damage from a spinal cord injury is often severe and irreversible, aspiration, incontinence and the other physical complications are problems that will not resolve with time.

Some patients who have a spinal cord injury may benefit from high-dose steroid therapy with drugs such as methylprednisolone. Using high-dose steroid therapy was once routine treatment for spinal cord

injuries, but in recent years the effectiveness of this therapy has been questioned. It is still used, but not as often and more selectively.

### **Transient Ischemic Attack**

A transient ischemic attack or TIA is defined as a sudden episode of neurological dysfunction that does not cause permanent damage. A TIA is similar to a stroke; it is caused by an interruption in blood flow to the brain, and the patients have many of the same signs and symptoms as do stroke patients. The causes of TIA and the factors that increase the risk of having a TIA are essentially the same as those of a stroke. Women are less likely than men to have a TIA.

The informal term for a TIA is a “mini-stroke” and, as noted above, there are similarities of a TIA to a stroke. However, there are two primary differences between a TIA and stroke that are important to know. In a TIA, the flow of blood is spontaneously restored and brain tissue is not damaged. A patient having a TIA will have many of the same signs and symptoms as someone having a stroke but the clinical presentation is usually less severe and the brain tissue will not be permanently damaged.

A TIA, however, is still a serious medical problem, and TIAs are very common. Approximately 20-30% of all people who have a TIA will have a stroke, and the stroke will happen 7 days or so after the TIA. Anyone who is having, or is suspected of having a TIA should be immediately transported to an emergency room.

### **Treatment of a TIA**

The primary goals when treating a patient who is having a TIA are:

1) make sure the patient is not having a stroke, 2) treat the medical

conditions that are causing the TIA, and 3) start therapies that will prevent the patient from having a stroke.

A computerized axial tomography (CAT) scan or a magnetic resonance imaging (MRI) scan, or some other type of imaging study should be done. This will determine if the patient is having a TIA or a stroke. The basic causes of the TIA such as atherosclerosis or hypertension will be treated, and some type of anticoagulant therapy will be started to prevent a stroke.

### **Cancers, Cardiac Arrest, Cardiac Arrhythmias, and Infections**

Cancers, cardiac arrest, cardiac arrhythmias, and infections can all cause damage to the brain and the spinal cord. Cancers and infections are not acute problems. These diseases tend to develop slowly, and the damage they cause to the brain and spinal cord is not sudden and dramatic. Cardiac arrest and cardiac arrhythmias such as ventricular tachycardia and ventricular fibrillation cause brain injury by an abrupt disruption of circulation. In these situations, the heart stops pumping or the pumping action of the heart is too weak and uncoordinated to deliver blood to the brain.

### **Patient Care: Stroke or Spinal Cord injury**

Caring for a patient who has had a stroke or a spinal cord injury is very involved. The basics of care should be that a CNA should particularly focus on include the following.

#### **Aspiration**

The health clinician or a therapist will prescribe specific aspiration prevention techniques such as elevation of the head while eating and swallowing exercises. The CNA must be aware that the signs of aspiration can be subtle, so any symptoms or unusual behaviors that seem to be associated with eating should be reported.

### **Nutrition**

Adequate nutrition and good hydration are essential for recovery. The CNA has a responsibility to monitor the patient's food and fluid intake.

### **Complications**

Complications that are associated with stroke and spinal cord injuries should be known, and the CNA should make sure to observe the patient closely for their development.

### **Skin Care**

Immobility, dehydration, incontinence, and pre-existing medical conditions such as diabetes are common in stroke and spinal cord injury patients. These increase the risk of developing skin breakdown and pressure ulcers, so the CNA must be very careful in the assessment and treatment of the patient's skin.

### **Case Study: Traumatic Brain Injury**

The following case study was obtained from a PubMed search and discusses the case of an elderly female diagnosed with a traumatic brain injury from bleeding within the brain.

The authors reported on a 79-year-old female who arrived to the hospital comatose from a traumatic brain injury. On physical

examination the patient was found to be deeply comatose. Laboratory showed normal results except for one test that is a marker for blood clot formation. A brain computed tomography (CT) revealed a large blood clot that first had to be surgically removed. She received the surgery to remove the brain blood clot. After surgery, there was mild improvement of her level of consciousness but the patient remained severely impaired. Another CT was done of the brain on the second day after surgery, which confirmed the area of injury to the brain. Treatment included blood thinner medication to treat the blood clot and shortly thereafter her level of consciousness improved and six months after surgery she was able to care for herself.

## **Discussion**

Blood clots to the brain from a large vein have always been considered to have poor outcomes. The common deep veins in people are more generally suspect when blood clots reach the brain. Causes known in the literature include pregnancy, oral contraceptives, infections, and brain cancers. More rare are a TBI or surgery that could lead to a blood clot to the brain.

The authors discussed the anatomy of the deep cerebral (brain) veins that drain blood from deep white brain matter and other deep structures of the brain. Patients suffering from a deep cerebral vein blood clot could present with headache, numbness, weakness, alteration of consciousness or other neuropsychological presentations. However, the authors also suggested that an accurate diagnosis of this type of brain blood clot tends to be difficult because of a low incidence of occurrence. Brain scans are helpful to diagnose this condition.

In this case, a combination of several techniques with clinical symptoms are recommended to diagnose the condition. A combination of laboratory testing showing a blood clotting issue and brain scans helped to diagnose the cause of the patient's TBI, which was a blood clot that caused injury in the deep area of the brain.

The authors identified first-line treatment to be anticoagulation or blood thinners, such as heparin. The patient's level of consciousness and recovery verified the efficacy of this treatment. More aggressive therapies are available such as surgical stenting, but this was not needed in this woman's case. The proper treatment involves prompt identification of the source of the blood clot, restoring blood flow and preventing serious complications. If the patient had not recovered as well as she did, then she would have been moved on to palliative therapy that may prolong life to a certain degree. Fortunately, for this patient she was able to recover enough to resume prior levels of self-care.

The authors reported what they believed to be a unique and rare finding of brain blood clot. They suggested that diagnosis of a deep vein blood clot in the brain should be suspected when patients do not recover from treatment for loss of consciousness secondary to brain injury.

## **Summary**

Brain and spinal cord injuries are common medical emergencies. Brain injuries associated with medical conditions are the third leading cause of death in the U.S., and they are the leading cause of disability. Most

brain injuries are a complication of widespread diseases such as atherosclerosis, diabetes, and hypertension.

The brain could be considered the command center of the body. The spinal cord helps to control some basic body functions and to act as the major “power conductor” of the information that the brain sends and receives. Two basic causes of brain injuries include trauma and acquired conditions, and the most common type of brain injury is a stroke. Strokes are one of the leading causes of death in the U.S. The immediate medical care of a stroke is critically important. Someone who is having a stroke needs evaluation and treatment as soon as possible. Aspiration can be a serious problem after a stroke. Loss of the gag reflex is common and puts the patient at risk for aspiration and pneumonia. The health clinician or a therapist will prescribe specific aspiration prevention techniques.

The CNA must be aware that the signs of aspiration can be subtle, so any symptoms or unusual behaviors that seem to be associated with eating should be reported. Immobility, dehydration, incontinence, and pre-existing medical conditions such as diabetes are common in stroke and spinal cord injury patients.